

Amendments to the Claims

Please amend the claims to read as follows:

1. (Withdrawn) A method of assessing relative susceptibility of a human to oxidative damage, the method comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least two genes selected from the group consisting of

- B3
- a) genes which encode an enzyme that catalyzes conversion of a toxic oxygen species to a less toxic oxygen species;
 - b) genes which encode a protein that provides protection against oxidative stress;
 - c) genes which encode a protein that induces production of a toxic oxygen species;
 - d) genes which encode a protein that indirectly affects oxidative stress; and
 - e) genes which encode a protein for which the level of expression of the protein is associated with oxidative stress,

whereby occurrence of any of the polymorphisms is an indication that the human is more susceptible to oxidative damage than a human whose genome does not comprise the polymorphism, and whereby occurrence of a plurality of the polymorphisms is an indication that the human is even more susceptible to oxidative damage than a human whose genome does not comprise the polymorphisms.

2-38. (Canceled)

39. (Amended) A method of selecting a dose of an anti-oxidant composition for administration to a human, the method comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least two genes selected from the group consisting of

- a) genes which encode an enzyme that catalyzes conversion of a toxic oxygen species to a less toxic oxygen species;
- b) genes which encode a protein that provides protection against oxidative stress;
and
- c) genes which encode a protein that induces production of a toxic oxygen species;
- d) ~~genes which encode a protein that indirectly affects oxidative stress; and~~

- e) ~~genes which encode a protein for which the level of expression of the protein is associated with oxidative stress;~~

whereby occurrence of any of the disorder-associated polymorphisms is an indication that a greater dose of the anti-oxidant composition should be administered to the human; and selecting a dose of the anti-oxidant composition based on occurrence of the disorder-associated polymorphisms.

40. (Withdrawn) A kit for assessing relative susceptibility of a human to oxidative damage, the kit comprising reagents for assessing occurrence in the human's genome of disorder-associated polymorphisms in at least two genes selected from the group consisting of

- B3
COT.
- a) genes which encode an enzyme that catalyzes conversion of a toxic oxygen species to a less toxic oxygen species;
 - b) genes which encode a protein that provides protection against oxidative stress;
 - c) genes which encode a protein that induces production of a toxic oxygen species;
 - d) genes which encode a protein that indirectly affects oxidative stress; and
 - e) genes which encode a protein for which the level of expression of the protein is associated with oxidative stress.

41-59. (Canceled)

60. (Previously Added) The method of claim 39, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least four genes selected from the group consisting of genes which encode an enzyme that catalyzes conversion of a toxic oxygen species to a less toxic oxygen species.

61. (Amended) The method of claim 60, wherein the four genes are

- i) the gene which encodes mitochondrial manganese superoxide dismutase (MnSOD),
- ii) the gene which encodes cytoplasmic copper/zinc superoxide dismutase (CZSOD),

- iii) the gene which encodes catalase, and
- iv) the gene which encodes glutathione peroxidase.

62. (Amended) The method of claim 39, ~~wherein the~~ comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least two genes ~~are~~ selected from the group consisting of

- i) the gene which encodes MnSOD,
- ii) the gene which encodes CZSOD,
- iii) the gene which encodes catalase,
- iv) the gene which encodes glutathione peroxidase,
- v) the gene which encodes glutathione S-transferase,
- vi) the gene which encodes glutathione reductase,
- vii) the gene which encodes thioredoxin reductase,
- viii) the gene which encodes paraoxonase,
- ix) the gene ~~genes~~ which encodes ~~encode~~ NAD(P)H:quinone oxidoreductase oxidoreductases 1 and 2,
- x) the gene which encodes 8-oxo-7,8-dihydrodeoxyguanosine triphosphatase,
- xi) the gene which encodes epoxide hydrolase,
- xii) the gene which encodes myeloperoxidase,
- xiii) the gene which encodes tumor necrosis factor alpha,
- xiv) the gene which encodes NADH/NADPH oxidase p22 phox protein,
- xv) the gene which encodes nitric oxide synthase,
- xvi) the gene which encodes xanthine oxidase,
- xvii) the gene which encodes cytochrome P450,
- xviii) the gene which encodes apolipoprotein E,
- xix) the gene which encodes UDP-glucuronosyltransferase 1A1,
- xx) the gene which encodes acid phosphatase,
- xxi) the gene which encodes protein phosphotyrosine phosphatase,
- xxii) the gene which encodes epinephrine oxidase,
- xxiii) the gene which encodes cystathionine beta-synthase,

B3
COD-4.

- xxiv) the gene which encodes cystathionine gamma-lyase,
- xxv) the gene which encodes N5-methyl THF:homocysteine methyltransferase,
- xxvi) genes which encode an S-adenosylmethionine methyltransferase, and
- xxvii) genes which encode a heat shock protein.

63. (Amended) The method of claim 62, ~~wherein the~~ comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least two genes are selected from the group consisting of i) through iv).

64. (Amended) The method of claim 62, ~~wherein the~~ comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least two genes are selected from the group consisting of i) through xi).

B3
COD1.
65. (Amended) The method of claim 62, ~~wherein the~~ comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least two genes are selected from the group consisting of i) through xvii).

66. (Previously Added) The method of claim 62, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least four of i) through xxvii).

67. (Previously Added) The method of claim 62, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least six of i) through xxvii).

68. (Previously Added) The method of claim 62, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least ten of i) through xxvii).

69. (Previously Added) The method of claim 62, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least fifteen of i) through xxvii).

70. (Amended) The method of claim 39, wherein occurrence of an individual disorder-associated polymorphism is assessed by

contacting a nucleic acid derived from the human's genome with a first oligonucleotide that anneals with higher stringency with a polynucleotide that consists of the disorder-associated polymorphism than with a polynucleotide that consists of a corresponding non-disorder-associated polymorphism and

assessing annealing of the first oligonucleotide and the nucleic acid,

whereby annealing of the first oligonucleotide and the nucleic acid is an indication that the human's genome comprises the disorder-associated polymorphism.

B3
CED 4.
71. (Previously Added) The method of claim 70, wherein the first oligonucleotide is attached to a support.

72. (Previously Added) The method of claim 71, wherein the support has a plurality of different first oligonucleotides attached thereto.

73. (Amended) The method of claim 72, wherein the support has attached thereto at least five first oligonucleotides that anneal with higher stringency with polynucleotides that consist of the disorder-associated polymorphisms than with polynucleotides that consist of the corresponding non-disorder-associated polymorphisms.

74. (Amended) The method of claim 72, wherein the support has attached thereto at least ten first oligonucleotides that anneal with higher stringency with polynucleotides that consist of the disorder-associated polymorphisms than with polynucleotides that consist of the corresponding non-disorder-associated polymorphisms.

75. (Amended) The method of claim 72, wherein the support has attached thereto at least fifteen first oligonucleotides that anneal with higher stringency with polynucleotides that consist of the disorder-associated polymorphisms than with polynucleotides that consist of the corresponding non-disorder-associated polymorphisms.

76. (Previously Added) The method of claim 70, wherein the first oligonucleotide is a molecular beacon oligonucleotide.

77. (Amended) The method of claim 70, wherein occurrence of an individual disorder-associated polymorphism is further assessed by

contacting the nucleic acid with a second oligonucleotide that anneals with higher stringency with a polynucleotide that consists of a non-disorder-associated polymorphism than with a polynucleotide that consists of the corresponding non-disorder-associated polymorphism and

assessing annealing of the second oligonucleotide and the nucleic acid,

whereby annealing of the second oligonucleotide and the nucleic acid is an indication that the human's genome does not comprise the disorder-associated polymorphism.

78. (Previously Added) The method of claim 77, wherein the second oligonucleotide is attached to a support.

79. (Previously Added) The method of claim 78, wherein the first and second oligonucleotides are attached to the same support.

80. (Previously Added) The method of claim 77, wherein the second oligonucleotide is a molecular beacon oligonucleotide.

81. (Previously Added) The method of claim 70, wherein the first and second oligonucleotides are spectrally distinct molecular beacon oligonucleotides.

82. (Previously Added) The method of claim 39, further comprising calculating a susceptibility score by summing, for each of the selected genes in which a disorder-associated polymorphism occurs in the human's genome, the product of a constant and a correlation factor, wherein the correlation factor represents the fraction of humans heterozygous or homozygous

for the disorder-associated polymorphism who exhibit the corresponding disorder, whereby the susceptibility score represents the relative susceptibility of the human to oxidative damage.

83. (Previously Added) The method of claim 82, wherein the same constant is used for each selected gene.

84. (Amended) The method of claim 82, wherein the constant used for each gene of group a) is greater than the constant used for the genes of groups b), ~~e)~~, d), ~~and e)~~ and c).

85. (Amended) The method of claim 84, wherein the constant used for each gene of group a) is at least twice as great as the constant used for the genes of groups b), ~~e)~~, d), ~~and e)~~ and c).

86. (Canceled).

87. (Amended) The method of claim ~~86~~ 85, wherein the genes are selected from the group consisting of

- i) the gene which encodes MnSOD,
- ii) the gene which encodes CZSOD,
- iii) the gene which encodes catalase,
- iv) the gene which encodes glutathione peroxidase,
- v) the gene which encodes glutathione S-transferase,
- vi) the gene which encodes glutathione reductase,
- vii) the gene which encodes thioredoxin reductase,
- viii) the gene which encodes paraoxonase,
- ix) the ~~gene~~ genes which ~~encodes~~ encode NAD(P)H:quinone ~~oxidoreductase~~ oxidoreductases 1 and 2,
- x) the gene which encodes 8-oxo-7,8-dihydrodeoxyguanosine triphosphatase,
- xi) the gene which encodes epoxide hydrolase,
- xii) the gene which encodes myeloperoxidase,
- xiii) the gene which encodes tumor necrosis factor alpha,

- xiv) the gene which encodes NADH/NADPH oxidase p22 phox protein,
- xv) the gene which encodes nitric oxide synthase,
- xvi) the gene which encodes xanthine oxidase, and
- xvii) the gene which encodes cytochrome P450.

88. (Previously Added) The method of claim 87, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least four of i) through xvii).

89. (Previously Added) The method of claim 87, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least six of i) through xvii).

B3
COD 4.
90. (Previously Added) The method of claim 87, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least ten of i) through xvii).

91. (Previously Added) The method of claim 87, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least fifteen of i) through xvii).

92. (Amended) The method of claim 39, wherein each of the disorder-associated polymorphisms is a single nucleotide polymorphism (SNP).

93. (Previously Added) The method of claim 92, wherein occurrence of a SNP is assessed by annealing a nucleic acid derived from the human's genome with a primer that is complementary to the region adjacent the SNP on its 3' side, extending the primer using a polymerase in order to add a nucleotide residue complementary to the SNP to the primer, and detecting the identity of the nucleotide residue complementary to the SNP.

94. (Previously Added) The method of claim 93, wherein the nucleotide residue is a non-extendable residue.

95. (New) The method of claim 39, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least two genes which encode an enzyme that catalyzes conversion of a toxic oxygen species to a less toxic oxygen species.

B3
cont.
96. (New) The method of claim 39, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least two genes which encode a protein that provides protection against oxidative stress.

97. (New) The method of claim 39, comprising assessing occurrence in the human's genome of disorder-associated polymorphisms in at least two genes which encode a protein that induces production of a toxic oxygen species
